arXiv:hep-th/9408042v1 8 Aug 1994

LPTENS-94/23
SWAT/93-94/38
hep–th/9408042
July 1994

LOTKA–VOLTERRA TYPE EQUATIONS AND THEIR EXPLICIT INTEGRATION

Jean-Loup GERVAIS
Laboratoire de Physique Théorique de l'École Normale Supérieure[1],
24 rue Lhomond, 75231 Paris CÉDEX 05, France.
and
Mikhail V. SAVELIEV[2]
Department of Mathematics, University College of Swansea,
Swansea SA2 8PP, Wales, U.K.

Abstract

In the present note we give an explicit integration of some two–dimensionalised Lotka–Volterra type equations associated with simple Lie algebras possessing a representation without branching.

---

[1] Unité Propre du Centre National de la Recherche Scientifique, associée à l'École Normale Supérieure et à l'Université de Paris-Sud.

[2] Permanent address: Institute for High Energy Physics, 142284, Protvino, Moscow region, Russia.

Equations of the Lotka–Volterra type under these or that boundary conditions appear in various branches of theoretical and mathematical physics, in particular in some solvable models of the field theory, plasma physics, electrotechnique, ecology, mathematical biology, etc. In the present note we discuss a special class of these equations associated with simple finite–dimensional Lie algebras. Namely, in paper [1] the finite system of the Lotka–Volterra type equations

$$\begin{aligned}\partial_+ R_{2\alpha-1} &= \lambda_\alpha e^{R_{2\alpha}} - \lambda_{\alpha-1} e^{R_{2\alpha-2}},\\ \partial_- R_{2\alpha} &= -e^{R_{2\alpha+1}} + e^{R_{2\alpha-1}},\ 1 \leq \alpha \leq N-1;\end{aligned} \tag{1}$$

was obtained for the functions $R_A(z_+, z_-)$, $1 \leq A \leq 2N$; with the boundary conditions $R_0 = R_{2N+1} = -\infty$. Here λ_α are some constants which can be eliminated, e.g. by redefinition of the functions $R_{2\alpha}$, only for the case related to the algebra A_r when none of them equal zero, see below. These equations realise, in fact, a relation between the Toda system

$$\partial_+ \partial_- x_i = \exp \rho_i; \quad \rho_i = \sum_{j=1}^{r} k_{ij} x_j,\ 1 \leq i \leq r; \tag{2}$$

associated with a complex simple Lie algebra $\mathcal{G}$ (of rank r with the Cartan matrix k) possessing a representation of dimension N without branching, and the $A_{N-2} \oplus \mathcal{G} \updownarrow (1)$–Toda fields. Seemingly, these equations can be extended to the system of 1st order partial differential equations associated with the relevant affine Lie algebras; that is rather important for constructing their soliton solutions.

Recall that speaking about a representation without branching, we mean such a representation of weight l, for which the representation space, endowed e.g. with the Verma base constructed with the help of the (cyclic) highest weight vector v_l by the action on it of the negative simple root vectors, $X_{-j_a} \cdots X_{-j_1} v_l$, $1 \leq j \leq r$, $1 \leq a \leq N-1$, contains the basis vectors $v_l(j_1, \cdots, j_a)$ with the only one value j_m corresponding to a given integer m. It is known that the 1st fundamental representations of A_r, B_r, C_r and G_2 possess this property.

Note that for the case of A_r system (1) realises the proper Bäcklund transformation, and, correspondingly, coincides with the two–dimensionalised difference KdV equations derived and completely integrated in [2] in terms of the general solutions to the A_r-Toda system, while in some different form than those of our formula (4) when $\mathcal{G} = A_r$.

Equations (1) represent an equivalent form of writing the system

$$\begin{aligned}\partial_-(e^{\theta_a} \Psi_a) &= e^{\theta_a} \Psi_{a+1}; & (3)\\ \partial_+ \Psi_a &= \lambda_a \exp \rho_{j_a} \Psi_{a-1}; & (4)\end{aligned}$$

$0 \leq a \leq N-1$, with the identification

$$R_{2\alpha-1} \equiv \log\, (\Psi_\alpha / \Psi_{\alpha-1}),\ R_{2\alpha} \equiv \rho_{j_\alpha} - \log\, (\Psi_\alpha / \Psi_{\alpha-1}).$$

Here we make use the following notations:

$$\lambda_a \equiv l_{j_a} - \sum_{q=1}^{a-1} k_{j_q j_a};\ \theta_a = \sum_j (l_j - \sum_{q=1}^{a} k_{j_q j}) x_j,\ \rho_{j_a} = \sum_j k_{j_a j} x_j.$$

When $\mathcal{G} = A_r$, equations (4) and (4), in terms of the functions $e^{\theta_{a-1}/2}\Psi_{a-1}$, are equivalent to the Frenet–Serret formulas for the moving frame [3]. For other simple Lie algebras we expect the similar situation with respect to the W–surfaces introduced in [4].

Due to their nilpotent structure, equations (4) and (4) can be solved in general form explicitly. Let us start from the first set of these equations, i.e. (4). (Recall that $\Psi_N \equiv 0$ by definition.) Taking $a = 0$, the corresponding equation is rewritten as

$$\partial_-(e^{\theta_0}\,\Psi_0) = e^{\theta_0}\,\Psi_1,$$

thereof

$$\Psi_1 \, = e^{-\theta_0}\partial_-(e^{\theta_0}\Psi_0),$$

where, in accordance with equation (4) for $a = 0$, the function Ψ_0 does not depend on z_+, $\Psi_0 = \Psi_0(z_-)$. Substituting this solution for Ψ_1 in the equation (4) for $a = 1$, i.e., in

$$\partial_-(e^{\theta_1}\,\Psi_1) = e^{\theta_1}\,\Psi_2,$$

we get

$$\Psi_2 \, = e^{-\theta_1}\partial_-(e^{\theta_1}\Psi_1) = e^{-\theta_1}\partial_-(e^{\theta_1-\theta_0}\partial_-(e^{\theta_0}\Psi_0)).$$

Continuing this integration procedure, we obtain the expressions for the whole set of the functions Ψ_a, $a > 0$, in terms of the function Ψ_0. The last step in the integration scheme, corresponding equation (4) with $a = N-1$, gives the equation of the N-th order for the function Ψ_0, the generalised Bargmann–Leznov generating equation [5]. This equation establish a relation between its coefficients (which are nothing but the W–potentials or conserved currents[3]) and the set of linearly independent fundamental solutions $\Psi_0^A(z_-)$, which are just the functions which determine, together with the analogous set of the functions $\Phi_0^A(z_+)$, the general solutions to the Toda system. The final result for the solutions R_A of system (1) can be written as:

$$e^{R_{2\alpha-1}} = \partial_- \,\log\,(e^{-\rho_{j_{\alpha-1}}}\partial_-(\cdots e^{-\rho_{j_1}}\partial_-\,(e^{\theta_0}\Psi_0))), \tag{5}$$

and $R_{2\alpha} = \rho_{j_\alpha} - R_{2\alpha-1}$. The corresponding expression for the functions Ψ_a contains a derivatives over z_-, and represents a generalisation of the determinant form of writing the formula for the moving frame for A_r given in [3]. These solutions, being substituted in equations (4), automatically satisfy them with account of the Toda system (2). The generating equation in the case under consideration here, can be written in the form

$$(\partial_z + \partial_z\theta_{N-1})\cdots(\partial_z + \partial_z\theta_1)\cdot(\partial_z + \partial_z\theta_0)\cdot\Psi_0(z) = 0. \tag{6}$$

This equation implies, in particular, the following expressions for the elements of the corresponding W–algebra:

$$W_p = \sum_{q_1=p}^{N}\sum_{q_2=p-1}^{q_1-1}\cdots\sum_{q_p=1}^{q_{p-1}-1}\mathcal{D}^0\cdot\mathcal{D}^1\cdots\mathcal{D}^{p-1}\cdot e^{\theta_{q_p-1}}, \tag{7}$$

---

[3] Note that a formulation of the elements of the corresponding W–algebra in terms of the Toda fields satisfying equations (2) comes back to the paper [6], see also [7]. At the same time, such differential polynomials in Toda fields arise also as the characteristic integrals providing the integrability property of these equations, see e.g. [8].

where $\mathcal{D}^p = \exp(\theta_{q_p-1} - \theta_{q_{p+1}-1}) \cdot \partial_z$.

The authors are grateful to L. N. Lipatov, D. I. Olive and A. V. Razumov for useful discussions. This work was partially supported by the Russian Fund for Fundamental Research, Grant # 94-01-01585-A; and ISF, Grant # RMO 000.

References

[1] M. V. Saveliev: Phys. Lett. A, **122** (1987) 312.

[2] A. N. Leznov, M. V. Saveliev, V. G. Smirnov: Lett. Math. Phys., **4** (1980) 445.

[3] Gervais, J.-L., Matsuo, Y.: Comm. Math. Phys., **152** (1993) 317.

[4] J.-L. Gervais, M. V. Saveliev: Preprint LPTENS-93/47, December 1993; Comm. Math. Phys. – to be published.

[5] A. N. Leznov: Lett. Math. Phys., **8** (1984) 5.

[6] A. Bilal, J.-L. Gervais: Phys. Lett., **B206** (1988) 412; Nucl. Phys., **B314** (1989) 646.

[7] J. Balog, L. Fehér, P. Forgacs, L. O'Raifeartaigh, A. Wipf: Ann. Phys., **203** (1990) 76.

[8] A. N. Leznov, M. V. Saveliev: *Group–Theoretical Methods for Integration of Nonlinear Dynamical Systems*, Birkhauser–Verlag, Progress in Physics, Basel, 1992.